



Propositional Logic, Truth Tables, and Predicate Logic (Rosen, Sections 1.1, 1.2, 1.3)

TOPICS

- Propositional Logic
- Logical Operations
- Equivalences
- Predicate Logic



Logic?





What is logic?

Logic is a truth-preserving system of inference

Truth-preserving:
If the initial statements are true, the inferred statements will be true

System: a set of mechanistic transformations, based on syntax alone

Inference: the process of deriving (inferring) new statements from old statements



Propositional Logic

- A *proposition* is a statement that is either true or false
- Examples:
 - This class is CS122 (true)
 - Today is Sunday (false)
 - It is currently raining in Singapore (???)
- Every proposition is true or false, but its *truth value* (true or false) may be unknown



Propositional Logic (II)

- A propositional statement is one of:
 - A simple proposition
 - denoted by a capital letter, e.g. 'A'.
 - A negation of a propositional statement
 - e.g. $\neg A$: "not A"
 - Two propositional statements joined by a *connective*
 - e.g. $A \wedge B$: "A and B"
 - e.g. $A \vee B$: "A or B"
 - If a connective joins complex statements, parenthesis are added
 - e.g. $A \wedge (B \vee C)$



Truth Tables

- The truth value of a compound propositional statement is determined by its truth table
- Truth tables define the truth value of a connective for every possible truth value of its terms



Logical negation

- Negation of proposition A is $\neg A$
 - A: It is snowing.
 - $\neg A$: It is not snowing
 - A: Newton knew Einstein.
 - $\neg A$: Newton did not know Einstein.
 - A: I am not registered for CS195.
 - $\neg A$: I am registered for CS195.



Negation Truth Table

A	$\neg A$
0	1
1	0



Logical and (*conjunction*)

- Conjunction of A and B is $A \wedge B$
 - A: CS160 teaches logic.
 - B: CS160 teaches Java.
 - $A \wedge B$: CS160 teaches logic and Java.
- Combining conjunction and negation
 - A: I like fish.
 - B: I like sushi.
 - I like fish but not sushi: $A \wedge \neg B$



Truth Table for Conjunction

A	B	$A \wedge B$
0	0	0
0	1	0
1	0	0
1	1	1



Logical or (*disjunction*)

- Disjunction of A and B is $A \vee B$
 - A: Today is Friday.
 - B: It is snowing.
 - $A \vee B$: Today is Friday or it is snowing.
- This statement is true if any of the following hold:
 - Today is Friday
 - It is snowing
 - Both
- Otherwise it is false



Truth Table for Disjunction

A	B	$A \vee B$
0	0	0
0	1	1
1	0	1
1	1	1



Exclusive Or

- The “or” connective \vee is inclusive: it is true if either *or both* arguments are true
- There is also an exclusive or \oplus

A	B	$A \oplus B$
0	0	0
0	1	1
1	0	1
1	1	0



Confusion over Inclusive OR and Exclusive OR

- Restaurants typically let you pick one (either soup or salad, not both) when they say “The entrée comes with a soup or salad”.
 - Use exclusive OR to write as a logic proposition
- Give two interpretations of the sentence using inclusive OR and exclusive OR:
 - Students who have taken calculus or intro to programming can take this class



Conditional & Biconditional Implication

- The conditional implication connective is \rightarrow
- The biconditional implication connective is \leftrightarrow
- These, too, are defined by truth tables

A	B	$A \rightarrow B$
0	0	1
0	1	1
1	0	0
1	1	1

A	B	$A \leftrightarrow B$
0	0	1
0	1	0
1	0	0
1	1	1



Conditional implication

- A: A programming homework is due.
- B: It is Tuesday.
- $A \rightarrow B$:
 - If a programming homework is due, then it must be Tuesday.
 - A programming homework is due only if it is Tuesday.
- Is this the same?
 - If it is Tuesday, then a programming homework is due.



Bi-conditional

- A: You can drive a car.
- B: You have a driver's license.
- $A \leftrightarrow B$
 - You can drive a car if and only if you have a driver's license (and vice versa).
- What if we said "if"?
- What if we said "only if"?



Compound Truth Tables

- Truth tables can also be used to determine the truth values of compound statements, such as $(A \vee B) \wedge (\neg A)$ (fill this as an exercise)

A	B	$\neg A$	$A \vee B$	$(A \vee B) \wedge (\neg A)$
0	0	1	0	0
0	1	1	1	1
1	0	0	1	0
1	1	0	1	0



Tautology and Contradiction

- A *tautology* is a compound proposition that is always true.
- A *contradiction* is a compound proposition that is always false.
- A *contingency* is neither a tautology nor a contradiction.
- A compound proposition is *satisfiable* if there is at least one assignment of truth values to the variables that makes the statement true.



Examples

A	$\neg A$	$A \vee \neg A$	$A \wedge \neg A$
0	1	1	0
1	0	1	0

Result is always true, no matter what A is.

Therefore, it is a **tautology**

Result is always false, no matter what A is.

Therefore, it is a **contradiction**



Logical Equivalence

- Two compound propositions, p and q , are logically equivalent if $p \leftrightarrow q$ is a tautology.
- Notation: $p \equiv q$
- De Morgan's Laws:
 - $\neg(p \wedge q) \equiv \neg p \vee \neg q$
 - $\neg(p \vee q) \equiv \neg p \wedge \neg q$
- How so? Let's build a truth table!



Prove $\neg(p \wedge q) \equiv \neg p \vee \neg q$

p	q	$\neg p$	$\neg q$	$(p \wedge q)$	$\neg(p \wedge q)$	$\neg p \vee \neg q$
0	0	1	1	0	1	1
0	1	1	0	0	1	1
1	0	0	1	0	1	1
1	1	0	0	1	0	0





Show $\neg(p \vee q) \equiv \neg p \wedge \neg q$

p	q	$\neg p$	$\neg q$	$(p \vee q)$	$\neg(p \vee q)$	$\neg p \wedge \neg q$
0	0	1	1	0	1	1
0	1	1	0	1	0	0
1	0	0	1	1	0	0
1	1	0	0	1	0	0



Other Equivalences

- Show $p \rightarrow q \equiv \neg p \vee q$
- Show Distributive Law:
 - $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$



Show $p \rightarrow q \equiv \neg p \vee q$

p	q	$\neg p$	$p \rightarrow q$	$\neg p \vee q$
0	0	1	1	1
0	1	1	1	1
1	0	0	0	0
1	1	0	1	1



Show $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$

p	q	r	$q \wedge r$	$p \vee q$	$p \vee r$	$p \vee (q \wedge r)$	$(p \vee q) \wedge (p \vee r)$
0	0	0	0	0	0	0	0
0	0	1	0	0	1	0	0
0	1	0	0	1	0	0	0
0	1	1	1	1	1	1	1
1	0	0	0	1	1	1	1
1	0	1	0	1	1	1	1
1	1	0	0	1	1	1	1
1	1	1	1	1	1	1	1





More Equivalences

Equivalence	Name
$p \wedge T \equiv p$ $p \vee F \equiv p$	Identity
$p \wedge q \equiv q \wedge p$ $p \vee q \equiv q \vee p$	Commutative
$p \vee (p \wedge q) \equiv p$ $p \wedge (p \vee q) \equiv p$	Absorption

See Rosen for more.



Equivalences with Conditionals and Biconditionals

- Conditionals
 - $p \rightarrow q \equiv \neg p \vee q$
 - $p \rightarrow q \equiv \neg q \rightarrow \neg p$
 - $\neg(p \rightarrow q) \equiv p \wedge \neg q$
- Biconditionals
 - $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$
 - $p \leftrightarrow q \equiv \neg p \leftrightarrow \neg q$
 - $\neg(p \leftrightarrow q) \equiv p \leftrightarrow \neg q$



Prove Biconditional Equivalence

p	q	$\neg q$	$p \leftrightarrow q$	$\neg(p \leftrightarrow q)$	$p \leftrightarrow \neg q$
0	0	1	1	0	0
0	1	0	0	1	1
1	0	1	0	1	1
1	1	0	1	0	0



Converse, Contrapositive, Inverse

- The *converse* of an implication $p \rightarrow q$ reverses the propositions: $q \rightarrow p$
- The *inverse* of an implication $p \rightarrow q$ inverts both propositions: $\neg p \rightarrow \neg q$
- The *contrapositive* of an implication $p \rightarrow q$ reverses and inverts: $\neg q \rightarrow \neg p$

The converse and inverse are not logically equivalent to the original implication, but the contrapositive is, and may be easier to prove.



Predicate Logic

- Some statements cannot be expressed in propositional logic, such as:
 - All men are mortal.
 - Some trees have needles.
 - $X > 3$.
- Predicate logic can express these statements and make inferences on them.



Statements in Predicate Logic

$P(x,y)$

- Two parts:
 - A predicate P describes a relation or property.
 - Variables (x,y) can take arbitrary values from some domain.
- Still have two truth values for statements (T and F)
- When we assign values to x and y , then P has a truth value.



Example

- Let $Q(x,y)$ denote “ $x=y+3$ ”.
 - What are truth values of:
 - $Q(1,2)$... false
 - $Q(3,0)$... true

- Let $R(x,y)$ denote x beats y in Rock/Paper/Scissors with 2 players with following rules:
 - Rock smashes scissors, Scissors cuts paper, Paper covers rock.
 - What are the truth values of:
 - $R(\text{rock}, \text{paper})$... false
 - $R(\text{scissors}, \text{paper})$... true



Quantifiers

- Quantification expresses the extent to which a predicate is true over a set of elements.
- Two forms:
 - Universal \forall
 - Existential \exists



Universal Quantifier

- $P(x)$ is true for all values in the domain
 $\forall x \in D, P(x)$
- For every x in D , $P(x)$ is true.
- An element x for which $P(x)$ is false is called a *counterexample*.
- Given $P(x)$ as “ $x+1 > x$ ” and the domain of \mathbb{R} , what is the truth value of:

$$\forall x P(x) \quad \dots \text{true}$$



Example

- Let $P(x)$ be that $x > 0$ and x is in domain of \mathbb{R} .
- Give a counterexample for:
 $\forall x P(x)$

$$x = -5$$




Existential Quantifier

- $P(x)$ is true for at least one value in the domain.

$$\exists x \in D, P(x)$$

- For some x in D , $P(x)$ is true.
- Let the domain of x be “animals”,
 $M(x)$ be “ x is a mammal” and
 $E(x)$ be “ x lays eggs”,
what is the truth value of:

$$\exists x (M(x) \wedge E(x))$$



English to Logic

- Some person in this class has visited the Grand Canyon.
- Domain of x is the set of all persons
- $C(x)$: x is a person in this class
- $V(x)$: x has visited the Grand Canyon
- $\exists x (C(x) \wedge V(x))$



English to Logic

- For every one there is someone to love.
- Domain of x and y is the set of all persons
- $L(x, y)$: x loves y
- $\forall x \exists y L(x, y)$
- Is it necessary to explicitly include that x and y must be different people (i.e. $x \neq y$)?
 - Just because x and y are different variable names doesn't mean that they can't take the same values



English to Logic

- No one in this class is wearing shorts and a ski parka.
- Domain of x is persons in this class
 - $S(x)$: x is wearing shorts
 - $P(x)$: x is wearing a ski parka
 - $\neg \exists x (S(x) \wedge P(x))$
- Domain of x is all persons
 - $C(x)$: x belongs to the class
 - $\neg \exists x (C(x) \wedge S(x) \wedge P(x))$



Evaluating Expressions: Precedence and Variable Bindings

- Precedence:
 - Quantifiers and negation are evaluated before operators
 - Otherwise left to right
- Bound:
 - Variables can be given specific values or
 - Can be constrained by quantifiers



Predicate Logic Equivalences

Statements are *logically equivalent* iff they have the same truth value under all possible bindings.

For example:

$$\forall x(P(x) \wedge Q(x)) \equiv \forall xP(x) \wedge \forall xQ(x)$$

In English: “Given the domain of students in CS160, all students have passed M124 course (P) and are registered at CSU (Q); hence, all students have passed M124 and all students are registered at CSU.



Other Equivalences

- Someone likes skiing (P) or likes swimming (Q); hence, there exists someone who likes skiing or there exists someone who likes swimming.

$$\exists x(P(x) \vee Q(x)) \equiv \exists xP(x) \vee \exists xQ(x)$$

- Not everyone likes to go to the dentist; hence there is someone who does not like to go to the dentist.

$$\neg \forall xP(x) \equiv \exists x\neg P(x)$$

- There does not exist someone who likes to go to the dentist; hence everyone does not like to go to the dentist.

$$\neg \exists xP(x) \equiv \forall x\neg P(x)$$